A lower bound on the distortion of embedding planar metrics into Euclidean space

Extended Abstract

Ilan Newman* Yuri Rabinovich*

Abstract

We exhibit a simple infinite family of series-parallel graphs which cannot be metrically embedded into Euclidean space with distortion smaller than $\Omega(\sqrt{\log n})$. This matches Rao's [15] general upper bound for metric embedding of planar graphs into Euclidean space, thus resolving the question how well do planar metrics embed in Euclidean spaces.

1 Introduction

Some of the most interesting questions in the study of finite metric spaces are about the relations between the structural properties of the underlying graph and of its geodetic metric (i.e., its shortest-path distance). In this paper we address one such question, and show a tight lower bound on the distortion of embedding a metric coming from a planar graph into Euclidean space.

Here are some basic definitions. A finite (semi-) metric space (S, μ) is a finite set S and a symmetric nonnegative distance function μ on $S \times S$ satisfying the triangle inequality and $\mu(x, x) = 0$. In what follows, we shall sometimes refer to μ as *metric*, regardless of S. A metric is called *planar* if it can be obtained by restricting the geodetic (i.e., shortest-path) metric of some weighted planar graph to a subset of its vertices. The weights should, of course, be nonnegative. *Series-parallel, tree*, etc., metrics are defined similarly. For example, one can easily check that the geodetic metric of the unit-weighted K_5 is a tree metric, while that of unit-weighted $K_{3,3}$ is not a planar metric.

Given two metric spaces (S, μ), (R, d) and an embedding $f : S \longrightarrow R$, the *distortion* of f is defined as

$$\mathrm{distr}(f) \;=\; \max_{x,y\in S} \frac{d(f(x), f(y))}{\mu(x, y)} \cdot \max_{x,y\in S} \frac{\mu(x, y)}{d(f(x), f(y))}\ ,$$

*CS Department, University of Haifa, Israel. Email: {ilan,yuri}@cs.haifa.ac.il. Supported in part by BSF grant 1999325.

i.e., the product of the maximum *expansion* and the maximum *contraction* of f. Observe that the distortion is never less than 1, and is equal to 1 exactly when f preserves μ up to scaling.

Two important parameters of a finite metric μ are $c_2(\mu)$ and $c_1(\mu)$, the smallest possible distortion of embedding μ into real Euclidean and ℓ_1 space respectively. Although the fundamental structural properties of finite metrics have just begun to emerge, numerous new exciting conjectures, theorems and applications (see, e.g., [8, 4, 10, 3, 12, 6, 15, 7]) seem to indicate that $c_1(\mu)$ and $c_2(\mu)$ indeed do capture some nontrivial aspects of μ.

Let us summarize the relevant facts about these two parameters. It always holds $c_2(\mu) \geq c_1(\mu)$, since any Euclidean metric is ℓ_1-embeddable [13]. Assuming μ is a metric on n points, both parameters are at most $O(\log n)$ [4], which is tight. The bound it is attained at the geodetic metric of any unit-weighted constant degree expander graph on n points [10, 1].

Tree metrics behave significantly better: $c_1(\mu)$ is 1, while $c_2(\mu) = O(\sqrt{\log\log n})$ [12], which is attained at the geodetic metric of the unit-weighted full binary tree of depth $\log n$ [4, 12]. A natural next question is what happens in the case when μ is a planar metric, or more generally, a metric coming from a graph excluding some fixed minor. An elegant result of Rao [15] shows that for such metrics $c_2(\mu)$, and consequently $c_1(\mu)$, are at most $O(\sqrt{\log n})$.

What are the true values of $c_1(\mu)$ and $c_2(\mu)$ for planar metrics? There are reasons to believe that $c_1(\mu)$ is constant; see [7] for a related discussion, and a proof that $c_1(\mu)$ is indeed constant for series-parallel and outerplanar graphs. For the last two years it was not clear whether $c_2(\mu)$ is closer to $O(\sqrt{\log n})$, as in the upper bound of [15], or to $O(\sqrt{\log\log n})$, as for trees [12].

In this paper we settle this question, and establish a lower bound of $\Omega(\sqrt{\log n})$ already for series-parallel metrics, which are a special case of planar metrics.

In addition to the main result, we obtain as a by-product of our construction an infinite family of weighted planar (in fact, tree-width 3) graphs H_n, so that not only the entire geodetic metric of H_n, but even the lengths of its edges must be distorted by a factor of $\Omega(\sqrt{\log n})$ by any embedding of H_n into Euclidean space. This contrasts with a construction of [11] showing that any weighted graph of tree-width 2 can be embedded into a line (!) so that the edges neither expand, nor contract by more than a factor of 3, and also with a theorem of Seymour [16] about the existence of edge-preserving metric embeddings of weighted planar graphs into ℓ_1 space.

2 The lower bound

We define a family $\{G_k\}$ of graphs in the following inductive manner, similar to the one used in [7]:

G_0 consists of a single edge. G_i is a "refinement" of G_{i-1} obtained by replacing each edge of G_{i-1} by two parallel paths, each containing two edges. The length of every edge in G_i is defined as 2^{-i}, half that of G_{i-1}. It is convenient to identify $V(G_{i-1})$ with the "old" vertices of G_i. Observe that this natural identification is in fact an isometry, i.e., the restriction of the geodetic metric of G_i to the "old" vertices is identical with the geodetic metric of G_{i-1} on these vertices.

Thus, without risk of confusion, we can speak of edges of G_i, and of their length, while discussing any G_k, $k \geq i$. To simplify the presentation, let us also introduce the notion of an anti-edge. Assume that the edge (a, b) of G_{i-1} was replaced in G_i by edges $(a, x), (x, b)$ and $(a, y), (y, b)$ respectively. The pair of vertices $\{x, y\} \subset V(G_i)$ will be called the *anti-edge* of (a, b). Observe that the distance between x and y is 2^{-i}, exactly as the length of the edge (a, b) of G_{i-1}.



Figure 2.1: The graph G_3.

It is easily checked that G_k is a series-parallel graph containing 4^k edges and $\frac{2 \cdot 4^k + 4}{3}$ vertices.

Theorem 2.1 *Let μ denote the geodetic metric of G_k. Then,*

$$c_2(\mu) \;\geq\; \sqrt{k+1}\;.$$

Proof: Let $f : V(G) \longrightarrow \mathbb{R}^d$ be an embedding of μ into Euclidean space. Due to the scalability of Euclidean space, we may w.l.o.g. assume that f is non-expanding, i.e., $\|f(v) - f(u)\|_2 \leq \mu(u, v)$ for any $u, v \in V(G_k)$. Let $\alpha = \min_{v,u \in V(G_k)} \frac{\|f(v)-f(u)\|_2}{\mu(v,u)}$. Our goal will be to show that $\alpha \;\leq\; 1/\sqrt{k+1}$, i.e., some distance must contract under such f by at least $\sqrt{k+1}$.

First, let us prove by downwards induction on i that the length of any edge of G_i, $i = k, k-1, ..., 0$, must contract under f by at least a factor of $[1 - (k-i)\alpha^2]^{-1/2}$. Formally, we claim that for any $(a, c) \in E(G_i)$,

$$\|f(a) - f(c)\|_2 \;\leq\; \sqrt{1 - (k-i)\alpha^2} \cdot \mu(a, c)\;.$$

The claim is trivially true for $i = k$, since f is non-expanding. Assume we have already demonstrated the claim for $i + 1$; let us demonstrate it for i. Recall the well known inequality that for any four points a, b, c, d in Euclidean space the sum of the squares of the diagonals never exceeds the sum of the squares of the sides:

$$\|a-b\|_2^2 \,+\, \|b-c\|_2^2 \,+\, \|c-d\|_2^2 \,+\, \|d-a\|_2^2 \;\geq\; \|a-c\|_2^2 \,+\, \|b-d\|_2^2\;. \tag{2.1}$$

The inequality holds since after subtracting the right hand side from the left hand side, one arrives at $\|a - b + c - d\|_2^2$.

Now, consider an edge (a, c) of G_i, its anti-edge (b, d), and the four surrounding edges of G_{i+1}: $(a, b), (b, c), (c, d)$ and (d, a). By our inductive assumption, the images of each of the latter pairs are at most $\sqrt{1 - (k-i-1)\alpha^2} \cdot 2^{-(i+1)}$ apart. By the definition of α, the images of b and d are at least $\alpha \cdot 2^{-i}$ apart. Combining this with (2.1), we get:

$$4 \cdot 2^{-2(i+1)} \cdot \left[1 - (k-i-1)\alpha^2\right] \;\geq\; \alpha^2 \cdot 2^{-2i} \;+\; \|f(a) - f(c)\|_2^2\;,$$

from which we conclude that

$$\|f(a) - f(c)\|_2^2 \ \leq \ \left[1 - (k-i)\alpha^2\right] \cdot 2^{-2i} \ = \ \left[1 - (k-i)\alpha^2\right] \cdot \mu^2(a,c) \ ,$$

as claimed.

Consider the edge (s,t) of G_0. The images of s and t are at least α apart by definition of α, and at most $\sqrt{1 - k \cdot \alpha^2}$ apart by the claim. Comparing the two terms we conclude that

$$\alpha \ \leq \ 1/\sqrt{k+1} \ .$$

■

Theorem 2.1 can be slightly strengthened. Observe that in our proof we have used only the edges and the anti-edges of G_i's. Therefore, restoring all these pairs as edges, and assigning them weight equal to their distance in G_k, we arrive at the graph H_k, whose edges must suffer distortion $\sqrt{k+1}$ in any embedding of H_k into Euclidean space. Formally:

Let H_0 consist of single edge of length 1, and let H_i be obtained by taking H_{i-1}, and in addition to existing vertices and edges, introducing for each edge $e = (a,c)$ of H_{i-1} of length $2^{-(i-1)}$ two new vertices b_e, d_e, a new edge (b_e, d_e) of length $2^{-(i-1)}$, and four new edges $(a, b_e), (b_e, c), (c, d_e)$ and (d_e, a), each of length 2^{-i}. As before, H_{i-1} isometrically embeds into H_i under the natural identification of the vertices.

It turns out that H_k is still planar, and, moreover, has tree-width 3. It has $\frac{5 \cdot 4^k - 2}{3}$ edges and the same number of vertices as in G_k. By the preceding discussion,

Theorem 2.2 *In any embedding f of H_k into Euclidean space which does not expand the edges, there exists an edge in $E(H_k)$ whose length is contracted by f by at least a factor of $\sqrt{k+1}$.*

We conclude this paper with a remark about a wider class of metric spaces for which one can prove a super-constant lower bound on the distortion of embedding of G_k. Assume that a metric space (R,d) has two properties, "stability" and "compactess". By stability we mean that for every $\delta > 0$ there exists an $\epsilon > 0$ such that if all the distances $d(a,x), d(x,b); d(a,y), d(y,b)$ are at most $0.5(1+\epsilon) \cdot d(a,b)$, then $d(x,y) \ \leq \ \delta \cdot d(a,b)$. Compactess means that the limit of a convergent sequence of (finite) (R,d)-embeddable metric spaces is also (R,d)-embeddable. The two properties imply that the distortion of embedding G_k into (R,d) tends to infinity with k. Indeed, it is not hard to see that the stability implies that the limit metric μ_∞ of G_∞ cannot be embedded into (R,d) with a bounded distortion. Applying the compactness argument, the conclusion follows. Examples of such metric spaces include e.g., ℓ_p spaces with $\infty > p > 1$, and hyperbolic spaces H^n of arbitrary dimension n.

Acknowledgments

We are grateful to Anupam Gupta for mentioning to us to the construction of Rao, and clarifying it. Thanks also to Nati Linial and Jirka Matoušek for insightful discussions.

References

[1] Yonatan Aumann and Yuval Rabani. An $O(\log k)$ approximate min-cut max-flow theorem and approximation algorithm. *SIAM Journal on Computing*, 27(1):291–301, February 1998.

[2] Yair Bartal. Probabilistic approximations of metric spaces and its algorithmic applications. In *Proceedings of the 37th Annual IEEE Symposium on Foundations of Computer Science*, pages 184–193, 1996.

[3] Yair Bartal. On approximating arbitrary metrics by tree metrics. In *Proceedings of the 30th Annual ACM Symposium on Theory of Computing*, pages 161–168, 1998.

[4] Jean Bourgain. On Lipschitz embeddings of finite metric spaces in Hilbert space. *Israel Journal of Mathematics*, 52(1-2):46–52, 1985.

[5] Jean Bourgain. The metrical interpretation of superreflexivity in Banach spaces. *Israel Journal of Mathematics*, 56(2):222–230, 1986.

[6] Uriel Feige. Approximating the bandwidth via volume respecting embeddings. In *Proceedings of the 30th Annual ACM Symposium on Theory of Computing*, pages 90–99, 1998.

[7] A. Gupta, I. Newman, Y. Rabinovich and A. Sinclair, *Cuts, trees and ℓ_1-embeddings of graphs.* In *Proceedings of the 40tn Annual Symposium on Foundatitions of CS*, pages 399–408, 1999.

[8] W.B. Johnson and J. Lindenstrauss, Extensions of Lipschitz mappings into a Hilbert space, *Contemporary Mathematics* 26 (1984), 189-206.

[9] Philip Klein, Serge A. Plotkin, and Satish B. Rao. Excluded minors, network decomposition, and multicommodity flow. In *Proceedings of the 25th Annual ACM Symposium on Theory of Computing*, pages 682–690, 1993.

[10] Nathan Linial, Eran London, and Yuri Rabinovich. The geometry of graphs and some of its algorithmic applications. *Combinatorica*, 15(2):215–245, 1995.

[11] Nathan Linial, Jiří Matoušek and Yuri Rabinovich. Embedding the edges of series-parallel graphs into line. Manuscript. 2000.

[12] Jiří Matoušek. On embedding trees into uniformly convex Banach spaces. *Israel Journal of Mathematics*, 114:221–237, 1999. (Czech version in : *Lipschitz distance of metric spaces*, C.Sc. degree thesis, Charles University, 1990).

[13] V. Milman, G. Schechtman. *Asymptomic Theory of Finite Dimensional Spaces*, LNM 1200, Springer-Verlag 1986.

[14] Yuri Rabinovich. Remarks on Klein-Plotkin-Rao Lemma. Manuscript.

[15] Satish B. Rao. Small distortion and volume preserving embeddings for planar and Euclidean metrics. In *15th Annual ACM Symposium on Computational Geometry*, pages 300–306, 1999.

[16] P. D. Seymour. Sums of circuits. In J.A. Bondy and U.S. Murty, editors, *Graph theory with applications*, American Elsevier, NY, 1979. pages 341–355.